

Charged and neutral binary nucleation of sulfuric acid in free troposphere conditions

Jonathan Duplissy, Joonas Merikanto, Karine Sellegri, Clemence Rose, Eija Asmi, Evelyn Freney, Heikki Juninen, Mikko Sipilä, Hanna Vehkamäki, Markku Kulmala, and CLOUD collaboration

Citation: [AIP Conference Proceedings](#) **1527**, 181 (2013); doi: 10.1063/1.4803233

View online: <http://dx.doi.org/10.1063/1.4803233>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1527?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Thermodynamic modeling of atmospheric aerosols: 0-100% relative humidity](#)

AIP Conf. Proc. **1527**, 441 (2013); 10.1063/1.4803299

[Multi-species nucleation rates in CLOUD](#)

AIP Conf. Proc. **1527**, 326 (2013); 10.1063/1.4803269

[Aerosol nucleation and growth in a mixture of sulfuric acid/alpha-pinene oxidation products at the CERN CLOUD chamber](#)

AIP Conf. Proc. **1527**, 322 (2013); 10.1063/1.4803268

[Molecular steps of neutral sulfuric acid and dimethylamine nucleation in CLOUD](#)

AIP Conf. Proc. **1527**, 302 (2013); 10.1063/1.4803263

[On atmospheric neutral and ion clusters observed in Hyytiälä spring 2011](#)

AIP Conf. Proc. **1527**, 169 (2013); 10.1063/1.4803230

Charged and Neutral Binary Nucleation of Sulfuric Acid in Free Troposphere Conditions

Jonathan Duplissy^a, Joonas Merikanto^a, Karine Sellegri^b, Clemence Rose^a, Eija Asmi^b, Evelyn Freney^b, Heikki Juninen^b, Mikko Sipilä^a, Hanna Vehkamäki^a, Markku Kulmala^a and the CLOUD collaboration

^a *University of Helsinki, Department of Physics, P.O. Box 64, 00014 University of Helsinki, Finland*

^b *Laboratoire de Météorologie Physique CNRS UMR6016, Observatoire de Physique du Globe de Clermont-Ferrand, Université Blaise Pascal, France*

Abstract. We present a data set of binary nucleation of sulfuric acid and water, measured in the CLOUD chamber at CERN during the CLOUD3 and CLOUD5 campaigns. Four parameters have been varied to cover neutral and ion-induced binary nucleation processes: Sulfuric acid concentration ($1\text{e}5$ to $1\text{e}8$ molecules per cm^{-3}), relative humidity (10% to 80%), temperature (208–293K) and ion concentration (0–4000 ions per cm^{-3}). In addition, classical nucleation theory implemented with hydrates and ion induced nucleation is compared with the data set. Our model and data are also compared with nucleation rates measured at Puy de Dome in the tropopause.

Keywords: binary nucleation, ion-induced nucleation, CLOUD, Free Troposphere.

PACS: 92.60.Mt, 82.60.Nh.

INTRODUCTION

Up to half of the aerosol acting as cloud condensation nuclei originates from aerosol nucleation[1]. The possible influence of cosmic rays on aerosol nucleation is of considerable interest[2]. The CLOUD experiment at CERN aims to study, under controlled conditions, nucleation processes as well as the effects of ions on nucleation. CLOUD was installed at CERN in 2009 and seven successful campaigns were carried out to study ion-induced and neutral binary nucleation of H_2SO_4 and H_2O , for tropospheric and stratospheric conditions. In this presentation, we will present the nucleation data obtained in the CLOUD chamber together with the new model which includes an ion-induced nucleation mechanism. To complete the picture, first molecular observation of nucleation in the tropopause will be shown and compared with the CLOUD data and the newly improved classical nucleation theory.

LABORATORY MEASUREMENT

Technical input for the CLOUD design was obtained in a pilot experiment in 2006 [3]. The chamber is a 3m-diameter electro-polished stainless-steel cylinder (26.1 m³). A field cage is installed inside the chamber to allow the removal of ions, when required. The contents of the chamber are irradiated by UV light in the range 250-400 nm. Experimental runs can be performed at stable temperatures between 40°C and -65°C.

The chamber is exposed to a 3.5 GeV/c secondary pion beam from the CERN PS, corresponding to the characteristic energies and ionization densities of cosmic ray muons in the lower troposphere. The beam intensity can be adjusted to cover the natural range from ground level to the stratosphere. Ultra-pure air is obtained from the evaporation of cryogenic liquid N₂ and liquid O₂. The air is humidified with a Nafion humidifier. Ozone is added to the air by UV irradiation. Trace gases such as SO₂ are added from gas cylinders containing pressurised N₂ as the carrier.

The chamber instrumentation includes PTRMS, CIMS, Nano-SMPS, CPC battery, PSM, API-ToF, NAIS, Gerdien, LOPAP, dew point sensor, SO₂ and O₃ analyser, as well as T, P and UV sensors. Sulphuric acid was generated by means of UV light, SO₂ and water.

From the API-ToF data, pure binary nucleation of sulphuric acid and water can be clearly identified[4]. Nucleation processes for species other than water and sulphuric acid have been excluded from this study.




FIGURE 1. An illustration of CLOUD in the T11 experimental zone at the CERN PS. The de-focused particle beam exits a dipole magnet (bottom right), crosses the aluminium hodoscope counter (middle) and then traverses the 3m-diameter CLOUD chamber, before being stopped by the concrete wall (top left). The instruments (blue boxes) analyse the contents of the chamber via sampling probes. The temperature of the air is precisely controlled and circulates between the chamber and the thermal housing. The clearing field for the removal of ions is shown by transparency (bottom and upper part of the chamber). A technical description can be found in the supplementary of Kirkby et al. [4].

MODEL

The classical nucleation theory including the hydrate interaction model and the ion-induced nucleation has been merged into one model. From this improved model an extension of the parameterisation from Vehkamäki et al. [5] has been developed, and compared with the CLOUD and Puy de Dome data.

FIELD MEASUREMENT

To complete this study, a field measurement at the high altitude research station, Puy de Dome has been performed during winter 2012 (Figure 2). The puy de Dôme research station is located at 1 465m above sea level in central France (45:460 N, 2:57 0E). A detailed description of the site is provided by Venzac et al. [6]. The Puy de Dôme research site contains a number of different instruments capable of resolving the physical, chemical, optical, and hygroscopic properties of aerosol particles. During this campaign nucleation processes have been measured with an API-ToF, NAIS and PSM. Comparison of these nucleation events with model and cloud data will be presented.



FIGURE 2. Unfreezing the inlets at the Puy de Dome research station after a night inside cloud.

ACKNOWLEDGMENTS

We would like to thank CERN for supporting CLOUD with important technical and financial resources, and for providing a particle beam from the CERN Proton

Synchrotron. This research has received funding from the EC Seventh Framework Programme (Marie Curie Initial Training Network "CLOUD-ITN" no. 215072, MC-ITN "CLOUD-TRAIN" no. 316662, and ERC-Advanced "ATMNUCLE" grant no. 227463), the German Federal Ministry of Education and Research (project nos. 01LK0902A and 01LK1222A), the Swiss National Science Foundation (project nos. 200020_135307 and 206620_130527), the Academy of Finland (Center of Excellence project no. 1118615), the Academy of Finland (135054, 133872, 251427, 139656, 139995, 137749, 141217, 141451), the Finnish Funding Agency for Technology and Innovation, the Nessling Foundation, the Austrian Science Fund (FWF; project no. P19546 and L593), the Portuguese Foundation for Science and Technology (project no. CERN/FP/116387/2010), the Swedish Research Council, Vetenskapsrådet (grant 2011-5120), the Presidium of the Russian Academy of Sciences and Russian Foundation for Basic Research (grants 08-02-91006-CERN and 12-02-91522-CERN), and the U.S. National Science Foundation (grants AGS1136479 and CHE1012293).

REFERENCES

1. Merikanto, J., Spracklen, D.V., Mann, G.W., Pickering, S.J., & Carslaw, K.S. *Atmospheric Chemistry and Physics*, 2009, 9, 8601–8616.
2. Carslaw, K. Harrison R. G. and Kirkby J., *Science*, 2002, Vol. 298 no. 5599 pp. 1732-1737 , doi: 10.1126/science.1076964
3. Duplissy, M. B. Enghoff, K. L. Aplin, F. Arnold, H. Aufmhoff, M. Avngaard, U. Baltensperger, T. Bando, R. Bingham, K. Carslaw, J. Curtius, A. David, B. Fastrup, S. Gagne, F. Hahn, R. G. Harrison, B. Kellett, J. Kirkby, M. Kulmala, L. Laakso, A. Laaksonen, E. Lillestol, M. Lockwood, J. Makela, V. Makhmutov, N. D. Marsh, T. Nieminen, A. Onnela, E. Pedersen, J. O. P. Pedersen, J. Polny, U. Reichl, J. H. Seinfeld, M. Sipila, Y. Stozhkov, F. Stratmann, H. Svensmark, J. Svensmark, R. Veenhof, B. Verheggen, Y. Viisanen, P. E. Wagner, G. Wehrle, E. Weingartner, H. Wex, M. Wilhelmsson, and P.M. Winkler, *Atmospheric Chemistry and Physics*, 10, 1635-1647, 2010, doi:10.5194/acp-10-1635-2010
4. Kirkby, J. Curtius, J. Almeida, E. Dunne, J. Duplissy, S. Ehrhart, A. Franchin, S. Gagne, L. Ickes, A. Kurten, A. Kupc, A. Metzger, F. Riccobono, L. Rondo, S. Schobesberger, G. Tsagkogeorgas, D. Wimmer, A. Amorim, F. Bianchi, M. Breitenlechner, A. David, J. Dommen, A. Downard, M. Ehn, R. C. Flagan, S. Haider, A. Hansel, D. Hauser, W. Jud, H. Junninen, F. Kreissl, A. Kvashin, A. Laaksonen, K. Lehtipalo, J. Lima, E. R. Lovejoy, V. Makhmutov, S. Mathot, J. Mikkila, P. Minginette, S. Mogo, T. Nieminen, A. Onnela, P. Pereira, T. Petaja, R. Schnitzhofer, J. H. Seinfeld, M. Sipila, Y. Stozhkov, F. Stratmann, A. Tome, J. Vanhanen, Y. Viisanen, A. Vrtala, P. E. Wagner, H. Walther, E. Weingartner, H. Wex, P. M. Winkler, K. S. Carslaw, D. R. Worsnop, U. Baltensperger, and M. Kulmala, *Nature*, 2011, 476 , 429-433, 2011 doi:10.1038/nature10343
5. Vehkamäki, H., M. Kulmala, I. Napari, K. E. J. Lehtinen, C. Timmreck, M. Noppel, and A. Laaksonen, *J. Geophys. Res.*, 2002, 107, 4622, doi:10.1029/2002JD002184.
6. Venzac, K. Sellegri, P. Villani, D. Picard, and P. Laj, *Atmospheric Chemistry and Physics*, 2009, 9, 1465-1478